

## Conference on 'Translating nutrition: integrating research, practice and policy' Postgraduate Symposium

### Brewers' spent grain; bioactivity of phenolic component, its role in animal nutrition and potential for incorporation in functional foods: a review

Aoife L. McCarthy<sup>1</sup>, Yvonne C. O'Callaghan<sup>1</sup>, Charles O. Piggott<sup>2</sup>, Richard J. FitzGerald<sup>2</sup> and Nora M. O'Brien<sup>1\*</sup>

<sup>1</sup>*School of Food and Nutritional Sciences, University College Cork, Cork, Republic of Ireland*

<sup>2</sup>*Department of Life Sciences, University of Limerick, Limerick, Republic of Ireland*

Brewers' spent grain (BSG) is a low-value co-product of the brewing industry produced in large quantities annually. This paper reviews the existing evidence regarding the phenolic component of BSG, focusing on composition, extraction and biofunctions such as antioxidant, anti-atherogenic, anti-inflammatory and anti-carcinogenic activities. Furthermore, the incorporation of BSG in foodstuffs will be discussed, including the use of BSG as an animal feed supplement and the potential of BSG to be incorporated into foods for human consumption. BSG contains hydroxycinnamic acids including ferulic acid, *p*-coumaric acid and caffeic acid; which have shown bioactivity in the pure form (antioxidant, anti-inflammatory, anti-atherogenic and anti-cancer). Phenolic extracts from BSG have also shown antioxidant potential, by protecting against oxidant-induced DNA damage, possibly by Fe chelation. Studies show that BSG has many benefits when used as a supplement in animal feed, such as increasing milk yield, milkfat content and in providing essential dietary amino acids. The incorporation of BSG in human foods such as cookies and ready-to-eat snacks has resulted in increased protein and fibre contents of the products, where the changes in organoleptic properties are controllable. It can be concluded that the phenolic component of BSG has potential bioactive effects, which are worth pursuing given that the inclusion of BSG into human foodstuffs is viable and beneficial.

#### **Brewers' spent grain: Phenolic acids: Animal nutrition: Functional foods: Bioactivity**

Brewers' spent grain (BSG) is the solid fraction of barley malt remaining after the production of wort. According to the Eurostat Data, BSG is the main by-product of the brewing industry, with approximately 3.4 million tonnes being produced annually in the EU<sup>(1)</sup>, at least 160 000 tonnes of which are produced in Ireland.

This solid residue contains water insoluble proteins in addition to the husk, pericarp and seed coat of the original barley grain<sup>(2)</sup>. Protein and fibre account for 20 and 70% of BSG dry matter, respectively, while the starch content of BSG is insignificant (due to the absence of starchy endosperm). Owing to its protein-rich composition, BSG

has the potential to be utilised in a manner similar to whey protein, providing health benefits for consumers. BSG is also rich in phenolic compounds, particularly ferulic acid and *p*-coumaric acid<sup>(3)</sup>, along with oligosaccharides and polysaccharides<sup>(4)</sup>. Emerging evidence, with regard to the ability of dietary phenolic compounds to exhibit anti-carcinogenic, anti-inflammatory and antioxidant activities<sup>(5,6)</sup> has led to significant interest in plant phenolic compounds particularly by the food industry, scientists and consumers.

To date, BSG has been widely used as an animal feed, particularly for cattle, to provide high amounts

**Abbreviations:** BSG, brewers' spent grain; COX-2, cyclooxygenase-isoform 2; DPPH, 1,1-diphenyl-2-picrylhydrazyl.

\*Corresponding author: Professor Nora M. O'Brien, fax +353 21 4270244, email nob@ucc.ie

**Table 1.** The approximate chemical composition of brewers' spent grain

Study	Component (as % dry matter*)						
	Protein	Lipid/Oil	Ash	Cellulose	Hemicellulose	Lignin	Starch
Beldman <i>et al.</i> <sup>(9)</sup>	23.8	–	3.5	15.1	24.8	–	2.0
Valverde <sup>(10)</sup>	24.0	6.0	–	17.0	39.0	4.0	–
Kanauchi <i>et al.</i> <sup>(11)</sup>	24.0	10.6	2.4	25.4	21.8	11.9	–
Santos <i>et al.</i> <sup>(12)</sup>	24.2	3.9	3.4	–	–	–	–
El-Shafey <sup>(13)</sup>	26.7	8.9	3.9	–	–	5.3	–
Mussatto & Roberto <sup>(14)</sup>	15.3	–	4.6	16.8	28.4	27.8	–
Xiros <sup>(15)</sup>	14.2	13.3	3.3	12.0	40.2	11.5	–
Treimo <sup>(16)</sup>	23.4	–	–	–	–	12.6	7.8


\*Values expressed as % dry matter, which has been documented in two studies to be 20.4<sup>(9)</sup> and 20%<sup>(13)</sup>.

of both protein and fibre. BSG is an excellent feed ingredient for ruminants, providing all the essential amino acids when combined with inexpensive N sources such as urea<sup>(7)</sup>. However, with the increased cost of disposal of the solid fraction, alternative uses are highly sought-after and it has been shown that BSG can be effectively integrated into ready-to-eat snacks to increase dietary fibre, crude protein and fat levels<sup>(1)</sup>. Other areas of successful research include the blending of BSG with flour for incorporation into cookies<sup>(8)</sup> and the addition of BSG to dough to improve the dietary fibre content in bread<sup>(1)</sup>.

This review details the existing evidence regarding BSG. A specific focus is placed on the potential bioactivities of phenolic compounds (particularly ferulic acid and *p*-coumaric acid) present in BSG, and the incorporation of BSG into foodstuffs, for both human and animal consumption.

### Composition of brewers' spent grain

Many studies have reported on the approximate composition of BSG, which contains protein, fat, cellulose, hemicellulose and lignin (Table 1). As shown, there is good consistency with regard to the composition of BSG. However, variations can arise due to differences in barley variety, harvesting time, characteristics of hops added and brewery technology<sup>(12)</sup>. BSG predominantly consists of the husk-pericarp-seed coat layers that are rich in cellulose, non-cellulosic polysaccharides, lignin, protein and fat. This is reflected in the composition of BSG (Table 1), and thus BSG can be regarded as a lignocellulosic material<sup>(4)</sup>. In addition to the components detailed in Table 1, it has been shown that BSG is also a valuable source of vitamins, minerals and amino acids, particularly for animal feeding. The vitamins present in BSG are biotin, folic acid, niacin, choline, riboflavin and thiamine, pantothenic acid and pyridoxine<sup>(7)</sup>. BSG is also reported to contain minerals such as Ca, Cu, Fe, Mn, K and Na<sup>(7,17)</sup> and both essential (including lysine, histidine, methionine, phenylalanine, tryptophan) and non-essential (including alanine, serine, glycine, proline) amino acids<sup>(7)</sup>. When combined with inexpensive N sources, such as urea, BSG can provide all the essential amino acids to ruminant animals<sup>(7)</sup>.



ferulic acid (C<sub>10</sub>H<sub>10</sub>O<sub>4</sub>)

R = OCH<sub>3</sub>

*p*-coumaric acid (C<sub>9</sub>H<sub>8</sub>O<sub>3</sub>)

R = H

**Fig. 1.** The general structure of hydroxycinnamic acid and the functional groups for ferulic and *p*-coumaric acids.

### Phenolic component of brewers' spent grain

#### *Phenolics present in brewers' spent grain*

Phenolic acids, particularly hydroxycinnamic acids and hydroxybenzoic acids are secondary plant metabolites found extensively in plant foods. Phenolic acids are currently the focus of much attention due to their potential to act as antioxidant, anti-inflammatory and anti-carcinogenic compounds<sup>(5,6)</sup>.

As previously mentioned, BSG consists predominantly of the husk-pericarp-seed coat and is largely made up of cell walls. Since most of the phenolic compounds of the barley grain are contained in the husk<sup>(4)</sup> and hydroxycinnamic acids accumulate in the cell walls, BSG is a potentially valuable source of phenolic acids.

There is evidence to suggest that ferulic acid and *p*-coumaric acid (as shown in Fig. 1) are present at relatively high concentrations in BSG<sup>(3)</sup>. Some of the existing literature regarding the presence of ferulic and *p*-coumaric acid is detailed in Table 2. Ferulic acid was found to be the most abundant hydroxycinnamic acid being present at concentrations ranging from 1860 to 1948 µg/g, while the *p*-coumaric levels ranged from 565 to 794 µg/g<sup>(23)</sup>. More recent evidence shows that BSG consists of 1.16% mono and dimeric phenolic acids, with 53% of the monomeric phenolic acids accounted for by ferulic acid. The vast majority of phenolic acids were also found to be in the bound form<sup>(20)</sup>. It has been reported that following ferulic and *p*-coumaric acids, the next most abundant phenolic acids in BSG were found to be sinapic, caffeic and

**Table 2.** The percentage (% dry weight) of bound phenolics, ferulic acid and *p*-coumaric acid present in brewers' spent grain

Study	Percentage (as % dry matter)		
	Bound phenolics	Ferulic acid	<i>p</i> -Coumaric acid
Bartolome <sup>(3)</sup>	–	~0.2	~0.1
Mandalari <sup>(18)</sup>	–	0.6	0.3
Athanasios <sup>(19)</sup>	–	0.2–0.3	0.1–0.2
Forsell <sup>(20)</sup>	–	0.5	0.5
Treimo <sup>(16)</sup>	0.7	–	–
Robertson <sup>(21)</sup>	0.8	–	–
Szwajgier <sup>(22)</sup>	–	0.3	0.1

syringic acids<sup>(22)</sup>. A summary of the phenolic acids present in BSG is given in Table 3.

#### *Extraction of phenolic acids from brewers' spent grain*

Numerous studies have been conducted to extract phenolic acids from BSG. Novel techniques for extraction, such as a rapid microwave-assisted derivatisation process have been investigated<sup>(19)</sup>. However, the majority of approaches use the basis of either acid hydrolysis or saponification (with 1–4 M NaOH) and liquid–liquid or liquid–solid extraction. A review of methods of extracting, separating and detecting phenolic acids in natural plant foods showed that the most frequently used methods involve acid hydrolysis and saponification<sup>(24)</sup>. Extraction usually entails the use of solvents such as methanol and ethyl acetate. TLC is extensively used for detecting phenolic acids due to its high sample throughput. However, using HPLC gives a greater degree of separation of compounds and is highly reproducible where quantification is possible. Therefore, reverse phase-HPLC is predominantly used, but HPLC coupled with UV or diode array detection is also an option<sup>(25)</sup>. A review looking at the extraction and quantification of phenolics in foods also reported that methanolic extraction and alkaline hydrolysis are commonly used for phenolic acid extraction, while a sequential alkaline hydrolysis releases bound phenolics<sup>(26)</sup>. A new method has recently been developed and validated, for the release of phenolic acids (both free and bound) from cereals including barley. This method uses solid-phase extraction coupled with HPLC-diode array detection analysis and is simple, inexpensive and gives good recoveries and precision<sup>(27)</sup>. Recently published results show that exogenous ferulic acid esterase produced by the probiotic organism *Lactobacillus acidophilus* K1 can successfully release the free phenolics from BSG<sup>(28)</sup>. In 2005, a study looking at the hydroxycinnamate content of BSG fractions utilised saponification with 4 M NaOH. The supernatants were then neutralised and extracted with ethyl acetate, dried and re-suspended in MeOH:H<sub>2</sub>O<sup>(18)</sup>. More recent research also used this method, giving comparable results<sup>(21)</sup>. Using a LUNA C18 reverse phase-HPLC column, both studies found that ferulic acid was the phenolic acid in greatest abundance in BSG, with coumaric acid being second highest. Saponification (involving the treatment of samples

**Table 3.** The most abundant phenolic acids present in brewers' spent grain modified from Szwajgier *et al.*<sup>(22)</sup>

Phenolic acid	Free acid concentration (mg/100 g dry matter)	
	Mean	SE
Ferulic acid	336.3	16.0
<i>p</i> -Coumaric acid	64.4	4.6
Sinapic acid	42.0	1.1
Caffeic acid	9.9	0.7
Syringic acid	6.5	0.1
4-OH-benzoic acid	1.2	0.6
Chlorogenic acid	0.6	0.2
Protocatechuic acid	0.5	0.1

with 1–4 M NaOH solution) has been widely used to extract hydroxycinnamic acids from BSG<sup>(3,23,29)</sup>.

Extraction methods similar to those used for BSG, have also been utilised with other materials such as wheat bran extracts<sup>(30)</sup> and apple waste extracts<sup>(31)</sup>.

#### *Potential health benefits of phenolic component of brewers' spent grain*

As previously mentioned, ferulic and *p*-coumaric acids are the phenolic acids at highest concentrations in BSG (Table 3). A lot of research has been conducted looking at the antioxidant activity of hydroxycinnamic acids, particularly ferulic and *p*-coumaric acids. A commonly used method for quantification of antioxidant activity is the 1,1-diphenyl-2-picrylhydrazyl (DPPH) assay, which measures the ability of the test compound to scavenge the DPPH radical. Chen and Ho<sup>(32)</sup> have shown the antioxidant potential of ferulic acid using the DPPH and Rancimat methods, but ferulic acid was a less potent antioxidant than caffeic acid and  $\alpha$ -tocopherol<sup>(33)</sup>. Caffeic acid has been shown to act as an antioxidant *in vitro* and scavenged radicals including DPPH and the superoxide anion<sup>(34)</sup>. It has also been shown, using the DPPH assay, that a number of hydroxycinnamic acids act as antioxidants, scavenging DPPH in the order caffeic acid > sinapic acid = ferulic acid > ferulates > *p*-coumaric acid<sup>(35)</sup>. Similarly, but using an alternative method, a study investigating the phenolic compounds in wheat bran extract and their antioxidant activity again found that ferulic acid was one of the strongest antioxidants using the  $\beta$ -carotene linoleic acid model system. The  $\beta$ -carotene linoleic acid model system assay is based on the principle that at a high temperature the oxidation of linoleic acid produces peroxides that decolourise  $\beta$ -carotene. The wheat bran extracts with highest ferulic acid concentrations (following alkaline hydrolysis) also exhibited higher antioxidant activity<sup>(30)</sup>. Ferulic acid and caffeic acid have been reported to have excellent antioxidant potential at low concentrations, with the ability to scavenge a range of free radicals. Both phenolic acids scavenge the reactive oxygen species and reactive nitrogen species, with concentration-dependent scavenging of NO, superoxide and 2,20-azino-bis-3-ethylbenzthiazoline-6-sulfonic acid radical. In line with the previously discussed evidence, caffeic acid was a stronger scavenger of the DPPH radical, but ferulic acid was better

at scavenging 2,20-azino-bis-3-ethylbenzthiazoline-6-sulfonic acid radical and NO<sup>(36)</sup>. In a recent study on beers, a direct correlation was found between ferric-reducing antioxidant power and a number of phenolic acids including ferulic, *p*-coumaric, caffeic, sinapic and vanillic acids<sup>(37)</sup>. A second study also showed that some of the phenolic acids present in beer correlate with the antioxidant activity measured by the DPPH radical and superoxide anion scavenging, metal chelation and reducing power; these include syringic and caffeic acids<sup>(38)</sup>. Since ferulic acid is so well recognised as an antioxidant, it is approved for use as a food additive in some countries to prevent oxidation<sup>(39,40)</sup>. In addition, it is important to note that while phenolic compounds can have an antioxidant effect, they have also been shown to act as pro-oxidants under certain conditions, thus inducing oxidative stress. Recent literature suggests that at low concentrations, many phenolics exhibit pro-oxidant behaviour, whereas the synthetic antioxidants, including  $\alpha$ -tocopherol, do not<sup>(41)</sup>. For caffeic acid and ferulic acid to act as pro-oxidants, higher concentrations are required<sup>(36)</sup>. It has also been found by using the comet assay that at high concentrations *p*-coumaric acid enhanced DNA breakage induced by H<sub>2</sub>O<sub>2</sub>. This may be due to the production of reactive oxygen species by *p*-coumaric acid as a result of its pro-oxidant activity<sup>(42)</sup>. It has been suggested that this pro-oxidant effect is related to the presence of metal ions in the body (for example due to tissue injury releasing Fe and Cu) and is of relevance for the bioactivity of phenolic compounds *in vivo*<sup>(43)</sup>. It is clear from a small number of *in vivo* studies that the hydroxycinnamic acids have antioxidant properties. Such studies are essential to understand the biological role of these phenolic acids<sup>(44)</sup>.

In addition to their antioxidant potential, there is increasing evidence to suggest that phenolic acids can have an anti-carcinogenic effect. Caffeic acid exhibits an anti-proliferative effect on several cancer cells including mammary gland, adenocarcinoma, lymphoblastic leukaemia<sup>(45)</sup> and cervical cancer cell lines as assessed using the MTT assay<sup>(45,46)</sup>. The cyclooxygenase-isoform 2 (COX-2) assay has been used for determination of the anti-cancer potential of these compounds. Overexpression of COX-2 increases the conversion of arachidonic acid to prostaglandins, which are important mediators of inflammation, and are associated with cancer. Phenolic acids including caffeic acid<sup>(47)</sup> and vanillic acid<sup>(48)</sup>, and polyphenols including epigallocatechin-3-gallate<sup>(49)</sup> and quercetin<sup>(50)</sup>, have been shown to inhibit the expression of COX-2, possibly reducing cancer risk. Apoptosis in cancer cell lines is also an indicator of anti-carcinogenic potential and can be assessed by a number of methods including DNA fragmentation and the Hoechst staining assay. Cinnamic acid derivatives induced apoptosis in human leukaemia (HL60) and colon cancer (SW480) cell lines, as measured by the aforementioned apoptosis methods<sup>(51)</sup>. In addition, the anti-apoptotic effect of phenolic compounds including ferulic acid and caffeic acid on human peripheral blood mononuclear cells was investigated<sup>(52)</sup>. Caffeic acid inhibited externalisation of phosphatidyl serine, which indicates the pre-apoptotic stages, and hence it was concluded to have an anti-apoptotic effect. DNA

fragmentation was analysed using an apoptotic DNA ladder kit. It was shown that pre-treating cells with caffeic acid, ferulic acid or ellagic acid before exposure to H<sub>2</sub>O<sub>2</sub> inhibited DNA fragmentation. Recently published data add to the evidence available in the area, by measuring the ability of phenolic compounds to modulate NF- $\kappa$ B activity. In the inflammatory process, NF- $\kappa$ B is a transcription factor, whose increased activation has been reported in several human cancers<sup>(53)</sup>. Free phenolic acids that can be found in cereal grains (including ferulic, caffeic, sinapic and *p*-coumaric acids) significantly modulate NF- $\kappa$ B activity in U9373 $\kappa$ B-LUC cells, with a desired level of modulation being achieved by the synergistic action of phenolic acids and other phenolic compounds<sup>(54)</sup>. Animal studies have also been carried out to determine the anti-carcinogenic potential of phenolic acids. An animal study to establish the effect of curcumin, chlorogenic acid, ferulic acid and caffeic acid on tumour promotion in the skin of mice showed that chlorogenic, ferulic and caffeic acid prevented the number of 12-*O*-tetradecanoylphorbol-13-acetate-induced tumours per mouse by 60, 28 and 35%, respectively<sup>(55)</sup>. Results of a later animal study suggest that ferulic acid not only inhibits the growth of aberrant crypt foci in the colon but also prevents the conversion of pre-neoplasia to malignant neoplasia<sup>(56)</sup>. A recently published review on plant phenolics reported that natural phenolics, including tea and fruit polyphenols, play an antagonistic role in all stages of cancer development and that further study on these compounds will provide information regarding their possible future pharmaceutical use<sup>(57)</sup>.

Cytokines are small cell signalling molecules involved in the inflammatory response, these include interleukins and interferons (for example interferon- $\gamma$ ). The ability of a compound to alter the production of a stimulated cytokine or NO indicates the compound's potential to act as an immune-modulator. Murakami *et al.*<sup>(58)</sup> investigated the effect of both ferulic acid (from rice bran) and FA15 (a derivative of ferulic acid) on NO synthase, COX-2 and TNF $\alpha$  in the RAW264-7 murine macrophage cell line. Unlike the ferulic acid isolated from rice bran, the synthesised FA15 derivative was found to inhibit the release of TNF $\alpha$  and reduce the protein expression of both nitric oxide synthase and COX-2<sup>(58)</sup>. Ferulic acid has also been shown to inhibit macrophage inflammatory protein-2 and TNF $\alpha$  production, induced by lipopolysaccharide in a macrophage cell line. The effect, although dose-dependent, was very weak compared with the effect of dexamethasone (a well-known inhibitor of interleukins)<sup>(59)</sup>. In Japanese Oriental medicines, *Cimicifuga heracleifolia* is often used as an anti-inflammatory drug. Ferulic acid has been shown to be among the main phenolic acids in *C. heracleifolia*<sup>(60)</sup>. Sakai *et al.*<sup>(61)</sup> showed that ferulic acid and isoferulic acid could reduce macrophage inflammatory protein-2 production in a dose-dependent manner in RAW264-7 cells. It was suggested that ferulic acid and isoferulic acid are responsible, at least in part, for the anti-inflammatory properties of the *C. heracleifolia* drug<sup>(61)</sup>. Recently published data have shown that ferulic acid and *p*-coumaric acid inhibited lipopolysaccharide-induced NO production and inducible NO synthase in macrophages<sup>(62)</sup>. This supports earlier evidence suggesting that these phenolic

**Table 4.** DNA damage in U937 cells following 24 h incubation with 0.5% (v/v) pale (P1–P4) or black (B1–B4) brewers' spent grain phenolic extracts or 0.1 µg/ml ferulic acid (adapted from McCarthy *et al.*<sup>(69)</sup>)

Sample	DNA damage (% tail DNA) oxidant – H <sub>2</sub> O <sub>2</sub>		DNA damage (% tail DNA) oxidant – SIN-1		DNA damage (% tail DNA) oxidant – 4-NQO		DNA damage (% tail DNA) oxidant – <i>t</i> -BOOH	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Control	3.2 <sup>†</sup>	0.5	4.3 <sup>†</sup>	0.4	3.1 <sup>†</sup>	0.3	4.7 <sup>†</sup>	0.6
Oxidant control	40.3 <sup>†</sup>	0.3	40.0 <sup>†</sup>	3.9	39.5 <sup>†</sup>	1.1	15.5 <sup>†</sup>	0.8
P1	30.3 <sup>†</sup>	7.3	32.8 <sup>†</sup>	4.9	39.0 <sup>†</sup>	6.8	14.5 <sup>†</sup>	0.8
P2	29.0 <sup>†</sup>	6.0	22.7 <sup>*†</sup>	1.9	40.7 <sup>†</sup>	4.3	22.6 <sup>†</sup>	3.0
P3	39.9 <sup>†</sup>	3.9	33.8 <sup>†</sup>	2.9	37.3 <sup>†</sup>	3.2	22.1 <sup>†</sup>	3.1
P4	23.1 <sup>†</sup>	3.2	32.2 <sup>†</sup>	6.1	37.8 <sup>†</sup>	3.6	17.4 <sup>†</sup>	2.6
B1	25.8 <sup>*†</sup>	7.5	33.6 <sup>†</sup>	2.9	36.1 <sup>†</sup>	5.7	13.6 <sup>†</sup>	0.9
B2	9.6 <sup>*†</sup>	1.1	6.9 <sup>*†</sup>	1.0	31.9 <sup>†</sup>	3.2	18.5 <sup>†</sup>	3.3
B3	16.0 <sup>*†</sup>	2.0	18.1 <sup>*†</sup>	0.3	33.4 <sup>†</sup>	2.4	16.6 <sup>†</sup>	1.8
B4	14.0 <sup>*†</sup>	3.8	24.1 <sup>*†</sup>	2.2	33.7 <sup>†</sup>	6.8	14.0 <sup>†</sup>	2.7
Ferulic acid	17.1 <sup>*†</sup>	0.8	30.8 <sup>†</sup>	3.0	32.6 <sup>†</sup>	5.2	11.0 <sup>†</sup>	1.9

SIN-1, 3-morpholinosydnonimine hydrochloride; 4-NQO, 4-nitroquinoline oxide; *t*-BOOH, *tert*-butylhydroperoxide.

<sup>†</sup>Data represent the mean of at least three independent experiments.

<sup>†</sup>Denotes a significant difference in DNA damage ( $P < 0.05$ ), relative to oxidant control. Statistical analysis by ANOVA followed by Dunnett's test.

acids can act as anti-inflammatory agents, by reducing TNF $\alpha$  induced IL-6 production in adipocytes. Quercetin, *p*-coumaric acid and resveratrol showed greatest inhibition of IL-6 production<sup>(63)</sup>.

Oxidised LDL is a well-recognised risk marker of CVD which is principally caused by atherosclerosis<sup>(64)</sup>. Evidence exists for the effect of hydroxycinnamic acids on the inhibition of LDL oxidation. Nardini *et al.*<sup>(65)</sup> demonstrated the antioxidant effect of hydroxycinnamic acid derivatives such as caffeic, ferulic and *p*-coumaric acids on LDL oxidation *in vitro*, with the use of the copper ion Cu<sup>2+</sup> as a catalyst. At a concentration of 100 µM, all phenolic acids except *p*-coumaric acid inhibited LDL oxidation; at 20 µM, ferulic acid inhibited about 92% of Cu-catalysed human LDL oxidation; at 5 µM only caffeic acid strongly inhibited the oxidation of LDL<sup>(65)</sup>. A second study using similar methodology also found that both ferulic and *p*-coumaric acid showed a dose-dependent inhibition of human LDL oxidation *in vitro* when tested at 5, 10 and 20 µM<sup>(66)</sup>. In agreement with these data, results of a later study showed sinapic, ferulic and *p*-coumaric acids inhibited LDL oxidation<sup>(67)</sup>. Other compounds that have shown the ability to reduce Cu-induced LDL oxidation include catechin<sup>(68)</sup>.

Our research group is the first, to our knowledge, to look specifically at phenolic extracts from BSG. Four extracts from pale BSG (P1, P2, P3 and P4) and four extracts from black BSG (where the grain is roasted to 200°C before brewing; B1, B2, B3 and B4) were analysed. Each extract results from a different step in the extraction process and hence contains different phenolic acid compositions. Extract 1 (P1, B1) contains free phenolics, extract 2 (P2, B2) contains bound phenolics, extract 3 (P3, B3) contains the remainder of bound phenolics and extract 4 (P4, B4) contains phenolics extracted with 110 mM NaOH<sup>(69)</sup>. The ability of the phenolic extracts to protect against oxidant-induced DNA damage was determined using the comet assay. In the U937 cell line, oxidative DNA damage was

induced by a range of oxidants; H<sub>2</sub>O<sub>2</sub>, 3-morpholinosydnonimine hydrochloride, 4-nitroquinoline oxide and *tert*-butylhydroperoxide. Table 4 shows the ability of the extracts to protect the cells against DNA damage. There was no protection against DNA damage induced by either 4-nitroquinoline oxide or *tert*-butylhydroperoxide. Ferulic acid and the black BSG extracts significantly reduced the DNA damage induced by H<sub>2</sub>O<sub>2</sub>, while P2, B2, B3 and B4 significantly reduced the percent tail DNA induced by 3-morpholinosydnonimine hydrochloride<sup>(69)</sup>. The four oxidants used have different mechanisms of action; damage induced by both H<sub>2</sub>O<sub>2</sub> and 3-morpholinosydnonimine hydrochloride involve the Fenton reaction which is an Fe-dependent reaction, 4-nitroquinoline oxide mimics the action of UV and Cu<sup>2+</sup> plays an essential role, whereas Fe does not<sup>(70)</sup>, *tert*-butylhydroperoxide causes lipid peroxidation and acts in a Ca<sup>2+</sup>-dependent manner and Fe plays less of a role than in H<sub>2</sub>O<sub>2</sub>-induced damage<sup>(71)</sup>. Therefore, BSG phenolic extracts may provide protection against oxidant-induced DNA damage by Fe chelation<sup>(69)</sup>.

In summary, there is increasing evidence to suggest that phenolic acids, including those found at highest concentrations in BSG, can confer potential health benefits including anti-inflammatory, antioxidant, anti-carcinogenic and anti-atherogenic effects. Recent data suggest that BSG has antioxidant potential and therefore further research on the phenolic compounds extracted from BSG is warranted.

### Incorporation of brewers' spent grain into feed/foodstuffs

#### *Animal food*

As previously mentioned, BSG contains approximately 20 and 70% protein and fibre, respectively, and it is due to this favourable chemical composition that it has great potential for use as a raw material/food ingredient<sup>(4)</sup>. BSG is an ingredient of significant importance for ruminants.

When administered with low-cost N sources such as urea, BSG can supply all the essential amino acids to the ruminants. The effect of BSG on milk yield and composition and the blood components of dairy cattle has also been studied<sup>(72)</sup>. The cattle received a diet consisting of ground maize, maize silage, soya bean meal and wheat bran, with the latter three being substituted with wet brewer's grain in the treatment group. The study showed the treatment group had an increased milk yield, milkfat and milk total solids content. Blood components such as glucose, cholesterol, Na and TAG were not affected. While the main outlet for BSG is currently as a feedstuff for dairy cattle, research has also been conducted looking at the benefits of BSG for use as a feed for poultry and fish. The effect of replacing rice bran in a fish diet with 10–40% brewery waste grains has been investigated<sup>(73)</sup>. The brewery waste used contained 19% crude protein, 18–20% crude fibre and had a good amino acid profile. It was found that carp (oily freshwater fish) had better growth performance on diets containing brewery waste than the control group. The authors attributed this enhanced growth performance to the high-quality protein contained in the waste grains. A more recent study showed that biodegraded BSG contained cysteine, lysine and methionine in addition to fourteen other amino acids<sup>(74)</sup>. Depending on the microbe used to degrade the BSG, different amino acid concentrations were found, with alanine consistently at highest concentrations. This composition was noted to be of particular importance for poultry as cysteine, lysine and methionine are the main amino acids required in poultry nutrition. For convenience, the possibility of producing dry BSG cakes suitable for long-term storage was examined using membrane filter press technology<sup>(13)</sup>. These dry cakes could be used as an animal feed at any time, or as a starting material for the production of other products using BSG.

In summary, evidence suggests that whole BSG, fed as part of a total mixed ration, has many nutritional benefits for a range of animals, particularly dairy cattle. This has resulted in the routine use of BSG as an animal feed for cattle.

#### *Human food*

In addition to its use as an animal feed, BSG has been incorporated into foodstuffs for human consumption. Given its low cost and high nutritional value, BSG makes an ideal ingredient for human foods such as biscuits and ready-to-eat snacks, particularly where there is a need to increase fibre content. In 1978, the possibility of preparing cookies with flour containing BSG at levels ranging from 5 to 60% was examined<sup>(8)</sup>. At 40% BSG addition, the physical qualities of the cookies were sustained. This supplementation level gave a 74% increase in N and increased crude fibre ten-fold. These results were supported by work published in 2002, where the authors looked at the effect of BSG (at levels of 5–25%) on the fibre content and quality of cookies<sup>(75)</sup>. As the level of addition of BSG increased, there was a significant increase in dietary fibre content. Another documented foodstuff suitable for the inclusion of BSG is ready-to-eat snacks<sup>(1)</sup>. BSG was added to the formulation mix (consisting of ingredients such as corn starch and wheat flour) at levels

ranging from 10 to 30%. The incorporation was successful, increasing dietary fibre and crude protein levels. Similarly, the addition of BSG into extruded snack food has been studied<sup>(76)</sup>. The maize flour of chickpea snacks was replaced with BSG at levels of 10, 20, 25 and 30%. The parameters measured included the effect of BSG supplementation on texture, colour, moisture, fat, fibre, starch, protein, phenolic compounds and antioxidant capacity. With increasing levels of BSG addition, the percent protein, fat and fibre content increased, while starch decreased. It was suggested that foods fortified with BSG be considered as functional foods. In a further study by Stojceska and Ainsworth<sup>(77)</sup>, BSG was incorporated at different levels (0–30%) into wheat flour breads treated with four different enzymes and the bread quality was then evaluated<sup>(77)</sup>. Similar to the previous study, it was found that the fibre content of the breads was significantly increased by BSG addition. The change in fat content was significantly linked to the addition of BSG. When addition of BSG is combined with the appropriate use of enzymes, the shelf life, texture and loaf volume can also be improved. Initially, it was thought that BSG was too granular for direct addition to food and that it would have to first be converted to flour before use. However, a study in 2009 demonstrated that BSG of various particle sizes could be effectively used in the production of frankfurters<sup>(78)</sup>. The control frankfurter had the highest score for acceptability, but the other products also had high scores, with the score decreasing with increase in particle size and reduction in fat levels. The authors suggested that BSG be used to produce low-fat high-fibre meat products. In addition to particle size, there are a number of points for consideration with the incorporation of BSG into foodstuffs. Firstly, there are concerns about appearance. When moist, BSG is brown in colour, thus it can only be effectively integrated into off-white products. Such foods include cookies and cakes. More importantly, it is imperative that the organoleptic properties of the foodstuff remain acceptable to consumers and are similar to the current commercially available products. The study by Prentice *et al.*<sup>(8)</sup>, demonstrated that BSG addition at a level of 15% was the upper limit for organoleptic acceptability<sup>(8)</sup>. At this level, the organoleptic quality was lowered but still remained acceptable to consumer panels. Similarly, Stojceska *et al.*<sup>(1)</sup> found that there was a limit to acceptability<sup>(1)</sup>. At a level of addition of 30%, physicochemical characteristics (such as texture, colour and hardness) remained acceptable. However, the authors concluded that addition of BSG at 20% level was optimal to maintain properties of commercially available snack foods. Where the protein hydrolysates are to be extracted and incorporated into foodstuffs, there is concern over the bitter taste of some peptides, due to the hydrophobic amino acid content<sup>(79)</sup>.

#### **Conclusion**

The literature shows that phenolic compounds including ferulic, caffeic and *p*-coumaric acid can have antioxidant, anti-cancer, anti-atherogenic and anti-inflammatory effects. Given that these phenolic acids are some of the major phenolics in BSG, it is expected that phenolic extracts

from spent grain may also exhibit similar properties and have the potential to be developed for a range of bioactivities. BSG currently functions as an animal feed, having many nutritional benefits. While some attempts have been made to incorporate the bioactive components of BSG into foodstuffs, further research in this area is needed. Given the potential bioactive nature of the phenolic extracts from BSG, and the large amounts of BSG produced annually as a low value co-product, it is imperative that an alternative use be explored.

### Acknowledgements

Funding for this research was provided under the National Development Plan, through the Food Institutional Research Measure, administered by the Department of Agriculture, Food and the Marine, Ireland. The authors declare no conflict of interest. A. L. M. wrote the manuscript with contributions from Y. C. O' C., C. O. P., R. J. F. and N. M. O'B.

### References

- Stojceska V, Ainsworth P & Plunkett A (2008) The recycling of brewer's processing by-product into ready-to-eat snacks using extrusion technology. *J Cereal Sci* **47**, 469–479.
- Townsley PM (1979) Preparation of commercial products from brewer's waste grain and trub. *MBAA Tech Q* **16**, 130–134.
- Bartolome B, Santos M, Jimenez J *et al.* (2002) Pentoses and hydroxycinnamic acids in brewer's spent grain. *J Cereal Sci* **36**, 51–58.
- Mussatto S, Dragone G & Roberto I (2006) Brewers' spent grain: generation, characteristics and potential applications. *J Cereal Sci* **43**, 1–14.
- Nagasaka R, Chotimarkorn C, Shafiqul IM *et al.* (2007) Anti-inflammatory effects of hydroxycinnamic acid derivatives. *Biochem Biophys Res Commun* **358**, 615–619.
- Yang CS, Landau JM, Huang MT *et al.* (2001) Inhibition of carcinogenesis by dietary polyphenolic compounds. *Annu Rev Nutr* **21**, 381–406.
- Huige, NJ (1994) Brewery by-products and effluents. In *Handbook of Brewing*, pp. 501–550 [WA Hardwick, editor]. New York: Marcel Dekker.
- Prentice N, Kissell L, Lindsay R *et al.* (1978) High-fiber cookies containing brewers' spent grain. *Cereal Chem* **55**, 712–721.
- Beldman G, Hennekam J & Voragen AGJ (1987) Enzymatic hydrolysis of beer brewers' spent grain and the influence of pretreatments. *Biotechnol Bioeng* **30**, 668–671.
- Valverde P (1994) Barley spent grain and its future. *Cerveza y Malta* **122**, 7–26.
- Kanauchi O, Mitsuyama K & Araki Y (2001) Development of a functional germinated barley foodstuff from brewer's spent grain for the treatment of ulcerative colitis. *J Am Soc Brew Chem* **59**, 59–62.
- Santos M, Jimenez J, Bartolome B *et al.* (2003) Variability of brewer's spent grain within a brewery. *Food Chem* **80**, 17–21.
- El-Shafey E, Gameiro M, Correia P *et al.* (2004) Dewatering of brewer's spent grain using a membrane filter press: A pilot plant study. *Sep Sci Technol* **39**, 3237–3261.
- Mussatto SI & Roberto IC (2005) Acid hydrolysis and fermentation of brewer's spent grain to produce xylitol. *J Sci Food Agric* **85**, 2453–2460.
- Xiros C, Topakas E, Katapodis P *et al.* (2008) Hydrolysis and fermentation of brewer's spent grain by *Neurospora crassa*. *Bioresour Technol* **99**, 5427–5435.
- Treimo J, Westereng B, Horn SJ *et al.* (2009) Enzymatic solubilization of brewers' spent grain by combined action of carbohydrases and peptidases. *J Agric Food Chem* **57**, 3316–3324.
- Pomeranz Y & Dikeman E (1976). From barley to beer – a mineral study. *Brew Dig* **51**, 30–32.
- Mandalari G, Faulds CB, Sancho AI *et al.* (2005) Fractionation and characterisation of arabinoxylans from brewers' spent grain and wheat bran. *J Cereal Sci* **42**, 205–212.
- Athanasios M, Georgios L & Michael K (2007) A rapid microwave-assisted derivatization process for the determination of phenolic acids in brewer's spent grains. *Food Chem* **102**, 606–611.
- Forssell P, Kontkanen H, Schols HA *et al.* (2008) Hydrolysis of brewers' spent grain by carbohydrate degrading enzymes. *J Inst Brew* **114**, 306–314.
- Robertson JA, I'Anson KJA, Treimo J *et al.* (2010) Profiling brewers' spent grain for composition and microbial ecology at the site of production. *LWT-Food Sci Technol* **43**, 890–896.
- Szwajgier D, Wako A, Targoski Z *et al.* (2010) The use of a novel ferulic acid esterase from *Lactobacillus acidophilus* K1 for the release of phenolic acids from brewer's spent grain. *J Inst Brew* **116**, 293–303.
- Hernanz D, Nuñez V, Sancho AI *et al.* (2001) Hydroxycinnamic acids and ferulic acid dehydromers in barley and processed barley. *J Agric Food Chem* **49**, 4884–4888.
- Stalikas CD (2007) Extraction, separation, and detection methods for phenolic acids and flavonoids. *J Sep Sci* **30**, 3268–3295.
- Zgórka G & Kawka S (2001) Application of conventional UV, photodiode array (PDA) and fluorescence (FL) detection to analysis of phenolic acids in plant material and pharmaceutical preparations. *J Pharmaceut Biomed* **24**, 1065–1072.
- Nacz M & Shahidi F (2004) Extraction and analysis of phenolics in food. *J Chromatogr A* **1054**, 95–111.
- Irakli MN, Samanidou VF, Biliaderis CG *et al.* (2012) Development and validation of an HPLC-method for determination of free and bound phenolic acids in cereals after solid-phase extraction. *Food Chem* **134**, 1624–1632.
- Szwajgier D (2011) The use of an extracellular ferulic acid esterase from *Lactobacillus acidophilus* K1 for the release of phenolic acids during mashing. *J Inst Brew* **117**, 427–434.
- Faulds C, Mandalari G, LoCurto R *et al.* (2004) Arabinoxylan and mono- and dimeric ferulic acid release from brewer's grain and wheat bran by feruloyl esterases and glycosyl hydrolases from *Humicola insolens*. *Appl Microbiol Biotechnol* **64**, 644–650.
- Kim KH, Tsao R, Yang R *et al.* (2006) Phenolic acid profiles and antioxidant activities of wheat bran extracts and the effect of hydrolysis conditions. *Food Chem* **95**, 466–473.
- McCann M, Gill C, O'Brien G *et al.* (2007) Anti-cancer properties of phenolics from apple waste on colon carcinogenesis in vitro. *Food Chem Toxicol* **45**, 1224–1230.
- Chen JH & Ho CT (1997) Antioxidant activities of caffeic acid and its related hydroxycinnamic acid compounds. *J Agric Food Chem* **45**, 2374–2378.
- Brand-Williams W, Cuvelier M & Berset C (1995) Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci Technol* **28**, 25–30.

34. Gulcin I (2006) Antioxidant activity of caffeic acid (3, 4-dihydroxycinnamic acid). *Toxicology* **217**, 213–220.
35. Kikuzaki H, Hisamoto M, Hirose K *et al.* (2002) Antioxidant properties of ferulic acid and its related compounds. *J Agric Food Chem* **50**, 2161–2168.
36. Maurya DK & Devasagayam TPA (2010) Antioxidant and prooxidant nature of hydroxycinnamic acid derivatives ferulic and caffeic acids. *Food Chem Toxicol* **48**, 3369–3373.
37. Piazzon A, Forte M & Nardini M (2010) Characterization of phenolics content and antioxidant activity of different beer types. *J Agric Food Chem* **58**, 10677–10683.
38. Zhao H, Chen W, Lu J *et al.* (2010) Phenolic profiles and antioxidant activities of commercial beers. *Food Chem* **119**, 1150–1158.
39. Graf E (1992) Antioxidant potential of ferulic acid. *Free Radical Biol Med* **13**, 435–448.
40. Itagaki S, Kurokawa T, Nakata C *et al.* (2009) In vitro and in vivo antioxidant properties of ferulic acid: a comparative study with other natural oxidation inhibitors. *Food Chem* **114**, 466–471.
41. Fukumoto L & Mazza G (2000) Assessing antioxidant and prooxidant activities of phenolic compounds. *J Agric Food Chem* **48**, 3597–3604.
42. Ferguson LR, Zhu S-T & Harris PJ (2005) Antioxidant and antigenotoxic effects of plant cell wall hydroxycinnamic acids in cultured HT-29 cells. *Mol Nutr Food Res* **49**, 585–593.
43. Morton LW, Caccetta RA-A, Puddey IB *et al.* (2000) Chemistry and biological effects of dietary phenolic compounds: relevance to cardiovascular disease. *Clin Exp Pharmacol Physiol* **27**, 152–159.
44. Shahidi F & Chandrasekara A (2010) Hydroxycinnamates and their *in vitro* and *in vivo* antioxidant activities. *Phytochem Rev* **9**, 147–170.
45. Gomes CA, Girão da Cruz T, Andrade JL *et al.* (2003) Anticancer activity of phenolic acids of natural or synthetic origin: a structure–activity study. *J Med Chem* **46**, 5395–5401.
46. Chang W-C, Hsieh C-H, Hsiao M-W *et al.* (2010) Caffeic acid induces apoptosis in human cervical cancer cells through the mitochondrial pathway. *Taiwan J Obstet Gynecol* **49**, 419–424.
47. Kang NJ, Lee KW, Shin BJ *et al.* (2009) Caffeic acid, a phenolic phytochemical in coffee, directly inhibits Fyn kinase activity and UVB-induced COX-2 expression. *Carcinogenesis* **30**, 321–330.
48. Kim MC, Kim SJ, Kim DS *et al.* (2011) Vanillic acid inhibits inflammatory mediators by suppressing NF- $\kappa$ B in lipopolysaccharide-stimulated mouse peritoneal macrophages. *Immunopharmacol Immunotoxicol* **33**, 525–532.
49. Hussain T, Gupta S, Adhami VM *et al.* (2005) Green tea constituent epigallocatechin 3 gallate selectively inhibits COX 2 without affecting COX 1 expression in human prostate carcinoma cells. *Int J Cancer* **113**, 660–669.
50. García-Mediavilla V, Crespo I, Collado PS *et al.* (2007) The anti-inflammatory flavones quercetin and kaempferol cause inhibition of inducible nitric oxide synthase, cyclooxygenase-2 and reactive C-protein, and down-regulation of the nuclear factor kappaB pathway in Chang Liver cells. *Eur J Pharmacol* **557**, 221–229.
51. Akao Y, Maruyama H, Matsumoto K *et al.* (2003) Cell growth inhibitory effect of cinnamic acid derivatives from propolis on human tumor cell lines. *Biol Pharm Bull* **26**, 1057–1059.
52. Khanduja KL, Avti PK, Kumar S *et al.* (2006) Anti-apoptotic activity of caffeic acid, ellagic acid and ferulic acid in normal human peripheral blood mononuclear cells: a Bcl-2 independent mechanism. *Biochim Biophys Acta* **1760**, 283–289.
53. Escárcaga RO, Fuentes-Alexandro S, García-Carrasco M *et al.* (2007) The transcription factor nuclear factor-kappa B and cancer. *Clin Oncol* **19**, 154–161.
54. Hole AS, Grimmer S, Jensen MR *et al.* (2012) Synergistic and suppressive effects of dietary phenolic acids and other phytochemicals from cereal extracts on nuclear factor kappa B activity. *Food Chem* **133**, 969–977.
55. Huang MT, Smart RC, Wong CQ *et al.* (1988) Inhibitory effect of curcumin, chlorogenic acid, caffeic acid, and ferulic acid on tumor promotion in mouse skin by 12-*O*-tetradecanoylphorbol-13-acetate. *Cancer Res* **48**, 5941–5946.
56. Kawabata K, Yamamoto T, Hara A *et al.* (2000) Modifying effects of ferulic acid on azoxymethane-induced colon carcinogenesis in F344 rats. *Cancer Lett* **157**, 15–21.
57. Dai J & Mumper RJ (2010) Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules* **15**, 7313–7352.
58. Murakami A, Nakamura Y, Koshimizu K *et al.* (2002) FA15, a hydrophobic derivative of ferulic acid, suppresses inflammatory responses and skin tumor promotion: comparison with ferulic acid. *Cancer Lett* **180**, 121–129.
59. Sakai S, Ochiai H, Nakajima K *et al.* (1997) Inhibitory effect of ferulic acid on macrophage inflammatory protein-2 production in a murine macrophage cell line, RAW264. 7. *Cytokine* **9**, 242–248.
60. He K, Pauli GF, Zheng B *et al.* (2006) Cimicifuga species identification by high performance liquid chromatography-photodiode array/mass spectrometric/evaporative light scattering detection for quality control of black cohosh products. *J Chromatogr A* **1112**, 241–254.
61. Sakai S, Kawamata H, Kogure T *et al.* (1999) Inhibitory effect of ferulic acid and isoferulic acid on the production of macrophage inflammatory protein-2 in response to respiratory syncytialvirus infection in RAW264-7 cells. *Mediat Inflamm* **8**, 173–175.
62. Kim EO, Min KJ, Kwon TK *et al.* (2012) Anti-inflammatory activity of hydroxycinnamic acid derivatives isolated from corn bran in lipopolysaccharide-stimulated Raw 264-7 macrophages. *Food Chem Toxicol* **50**, 1309–1316.
63. Yen G-C, Chen Y-C, Chang W-T *et al.* (2010) Effects of polyphenolic compounds on tumor necrosis factor- $\alpha$  (TNF- $\alpha$ )-induced changes of adipokines and oxidative stress in 3T3-L1 adipocytes. *J Agric Food Chem* **59**, 546–551.
64. Yoshida H & Kisugi R (2010) Mechanisms of LDL oxidation. *Clin Chim Acta* **411**, 1875–1882.
65. Nardini M, D'Aquino M, Tomassi G *et al.* (1995) Inhibition of human low-density lipoprotein oxidation by caffeic acid and other hydroxycinnamic acid derivatives. *Free Radical Biol Med* **19**, 541–552.
66. Meyer AS, Donovan JL, Pearson DA *et al.* (1998) Fruit hydroxycinnamic acids inhibit human low-density lipoprotein oxidation *in vitro*. *J Agric Food Chem* **46**, 1783–1787.
67. Andreassen MF, Landbo A-K, Christensen LP *et al.* (2001) Antioxidant effects of phenolic rye (*Secale cereale* L.) extracts, monomeric hydroxycinnamates, and ferulic acid dehydromers on human low-density lipoproteins. *J Agric Food Chem* **49**, 4090–4096.
68. Chen YC, Lin JT, Liu SC *et al.* (2011) Composition of flavonoids and phenolic acids in Lychee (*Litchi Chinensis*) flower extracts and their antioxidant capacities estimated with human LDL, erythrocyte, and blood models. *J Food Sci* **76**, C724–C728.
69. McCarthy AL, O'Callaghan YC, Connolly A *et al.* (2012) Phenolic extracts of brewers' spent grain (BSG) as functional ingredients—assessment of their DNA protective



- effect against oxidant-induced DNA single strand breaks in U937 cells. *Food Chem* **134**, 641–646.
70. Yamamoto K, Inoue S & Kawanishi S (1993) Site-specific DNA damage and 8-hydroxydeoxyguanosine formation by hydroxylamine and 4-hydroxyaminoquinoline 1-oxide in the presence of Cu(II): Role of active oxygen species. *Carcinogenesis* **14**, 1397–1404.
71. Kruszewski M, Iwaneńko T, Bartłomiejczyk T *et al.* (2008) Hepatic iron content corresponds with the susceptibility of lymphocytes to oxidative stress in neonatal pigs. *Mutat Res* **657**, 146–149.
72. Belibasakis N & Tsirgogianni D (1996) Effects of wet brewers grains on milk yield, milk composition and blood components of dairy cows in hot weather. *Anim Feed Sci Technol* **57**, 175–181.
73. Kaur V & Saxena P (2004) Incorporation of brewery waste in supplementary feed and its impact on growth in some carps. *Bioresour Technol* **91**, 101–104.
74. Essien J & Udotong I (2008) Amino acid profile of biodegraded brewers spent grains (BSG). *J Appl Sci Environ Manage* **12**, 109–111.
75. Öztürk S (2002). Effects of brewers' spent grains on the quality and dietary fibre content of cookies. *J Inst Brew* **108**, 23–27.
76. Ainsworth P, Ibanoglu S, Plunkett A *et al.* (2007) Effect of brewers spent grain addition and screw speed on the selected physical and nutritional properties of an extruded snack. *J Food Eng* **81**, 702–709.
77. Stojceska V & Ainsworth P (2008) The effect of different enzymes on the quality of high-fibre enriched brewer's spent grain breads. *Food Chem* **110**, 865–872.
78. Özvural EB, Vural H, Gökbulut *et al.* (2009) Utilization of brewer's spent grain in the production of frankfurters. *Int J Food Sci Technol* **44**, 1093–1099.
79. Clemente A (2000) Enzymatic protein hydrolysates in human nutrition. *Trends Food Sci Technol* **11**, 254–262.